

Study of Molecule-Metal Interfaces by Means of the Normal Incidence X-ray Standing Wave Technique

Giuseppe Mercurio

Forschungszentrum Jülich GmbH
Peter Grünberg Institute (PGI)
Functional Nanostructures at Surfaces (PGI-3)

Study of Molecule-Metal Interfaces by Means of the Normal Incidence X-ray Standing Wave Technique

Giuseppe Mercurio

Schriften des Forschungszentrums Jülich
Reihe Schlüsseltechnologien / Key Technologies

Band / Volume 49

ISSN 1866-1807

ISBN 978-3-89336-816-7

Contents

Acknowledgements	IV
Abstract	VI
1 Introduction	1
2 The normal incidence x-ray standing wave technique	5
2.1 Fundamentals of XSW	5
2.2 The standing wave	7
2.3 XSW detection: photoelectrons	9
2.4 Basics of an XSW experiment	10
2.5 Coherent position and coherent fraction	13
2.6 Experimental set-up	16
3 Torricelli	19
3.1 Introduction	19
3.2 Preparing reflectivity and electron yield profiles for fitting	21
3.3 Fitting the reflectivity	22
3.3.1 The fitting function	22
3.3.2 Calculation of the ideal reflectivity R_{theo} and phase ϕ_{theo}	25
3.3.3 Calculation of the structure factors	29
3.4 Fitting the electron yield profile	38
3.4.1 Fitting the coherent position and fraction	38
3.4.2 Calculation of the nondipolar correction parameters	40
3.4.3 Fitting the asymmetry parameter Q	45
3.5 Error analysis of the coherent position and fraction	48
3.6 Overview and conclusions	52


4	PTCDA on Ag(110)	55
4.1	Introduction	55
4.2	Experimental details	57
4.2.1	PTCDA deposition and coverage estimation	57
4.2.2	XPS acquisition parameters of NIXSW experiments	60
4.2.3	XPS background	62
4.2.4	Line shapes of the fitting components	63
4.3	Experimental results	66
4.3.1	Carbon	66
4.3.1.1	Two-component model	66
4.3.1.2	Multicomponent model	71
4.3.2	Oxygen	82
4.4	Discussion	88
4.4.1	Interpretation of PTCDA coherent fraction	88
4.4.2	Adsorption geometry of PTCDA on Ag(110)	91
4.5	Conclusions	99
5	K-doped PTCDA on Ag(110)	101
5.1	Introduction	101
5.2	NIXSW study	102
5.2.1	Experimental details	102
5.2.1.1	Potassium deposition and coverage estimation	102
5.2.1.2	XPS acquisition parameters of NIXSW experiments	104
5.2.1.3	XPS background	106
5.2.1.4	Line shapes of the fitting components	108
5.2.2	Experimental results	109
5.2.2.1	Carbon	109
5.2.2.2	Oxygen	114
5.2.2.3	Potassium	118
5.2.2.4	Silver	120
5.2.3	Discussion of NIXSW results	122
5.3	LEED study	126
5.3.1	Experimental results	126
5.3.1.1	Role of K amount	129
5.3.1.2	Additional diffraction spots	131
5.3.1.3	Role of annealing step	133
5.3.2	Discussion of LEED results	134

5.4	STM study	136
5.4.1	Experimental details	136
5.4.2	<i>Low</i> K coverage results	137
5.4.2.1	Overview	137
5.4.2.2	X phase	137
5.4.3	<i>Intermediate</i> K coverage results: stripe phase	139
5.4.3.1	Introduction	139
5.4.3.2	Real space and reciprocal space images	140
5.4.3.3	Vertical properties of the stripe phase	145
5.4.3.4	Lateral properties of the stripe phase	148
5.4.3.5	Potassium fingerprint	149
5.4.4	Summary of STM results	153
5.5	UPS and XPS study	155
5.5.1	Experimental details	155
5.5.2	Work function results	157
5.5.3	XPS results	159
5.5.4	Discussion of UPS and XPS results	160
5.6	Structural model	167
5.6.1	Description of the structural model	167
5.6.2	Discussion of the structural model	169
5.7	Conclusions	177
6	Molecular switches on Ag(111)	179
6.1	Introduction	179
6.2	Experimental details	181
6.2.1	Sample preparation	181
6.2.2	XPS acquisition parameters of NIXSW experiments	183
6.2.3	XPS background	184
6.2.4	Line shapes of the fitting components	185
6.3	Experimental results	186
6.3.1	Introduction	186
6.3.2	Silver	186
6.3.2.1	Ag3 <i>d</i>	187
6.3.2.2	Ag3 <i>d</i> plasmons	189
6.3.3	Azobenzene	193
6.3.3.1	Nitrogen	193
6.3.3.2	Carbon	200

Contents

6.3.4	TBA	202
6.3.4.1	Nitrogen	202
6.3.4.2	Carbon	209
6.4	Discussion	211
6.4.1	Discussion of azobenzene experimental results	211
6.4.1.1	Introduction	211
6.4.1.2	A novel interpretative scheme of NIXSW results	212
6.4.1.3	NIXSW adsorption model of azobenzene/Ag(111)	224
6.4.1.4	Comparison of NIXSW and DFT adsorption models	226
6.4.2	Discussion of TBA experimental results	230
6.4.2.1	Introduction	230
6.4.2.2	Comparison of NIXSW results with DFT calculations	231
6.4.2.3	NIXSW simulations of TBA/Ag(111)	232
6.4.2.4	NIXSW adsorption model of TBA/Ag(111)	240
6.5	Conclusions	245
7	Terephthalic acid on Cu(100)	249
7.1	Introduction	249
7.2	Experimental details	252
7.2.1	Sample preparation and coverage estimation	252
7.2.2	XPS acquisition parameters of NIXSW experiments	255
7.2.3	XPS background	256
7.2.4	Line shapes of the fitting components	257
7.3	Experimental results	258
7.3.1	Carbon	258
7.3.2	Oxygen	266
7.3.3	Copper	269
7.4	Discussion	273
7.4.1	Adsorption geometry of deprotonated TPA on Cu(100)	273
7.4.2	Comparison of NIXSW data with DFT calculations	282
7.5	Conclusions	294
8	Summary and conclusions	297
A	Technical details and code of Torricelli	301
A.1	Introduction	301
A.2	Preparing reflectivity and electron yield profiles for fitting	302
A.2.1	Technical details	302

A.2.2	Code	304
A.3	Fitting the reflectivity	306
A.3.1	Technical details	306
A.3.2	Code	307
A.4	Calculating the ideal reflectivity	310
A.4.1	Technical details	310
A.4.2	Code	311
A.5	Calculating the structure factors	318
A.5.1	Technical details	318
A.5.2	Code	320
A.6	Fitting the electron yield profile: coherent position and fraction . . .	325
A.6.1	Technical details	325
A.6.2	Code	327
A.7	Calculating the nondipolar correction parameters	331
A.7.1	Technical details	331
A.7.2	Code	332
A.8	Fitting the electron yield profile: the asymmetry parameter Q	334
A.8.1	Technical details	334
A.8.2	Code	335
Bibliography		355



Schlüsseltechnologien / Key Technologies
Band / Volume 49
ISBN 978-3-89336-816-7