



## Experimental determination of the partitioning coefficient of nopinone as a marker substance in organic aerosol

Bettina Steitz

Forschungszentrum Jülich GmbH  
Institute for Energy and Climate Research (IEK)  
Troposphere (IEK-8)

# **Experimental determination of the partitioning coefficient of nopinone as a marker substance in organic aerosol**

Bettina Steitz

Schriften des Forschungszentrums Jülich  
Reihe Energie & Umwelt / Energy & Environment

Band / Volume 169

---

ISSN 1866-1793

ISBN 978-3-89336-862-4

# Contents

<b>1</b>	<b>Introduction</b>	<b>11</b>
1.1	Atmospheric Aerosols . . . . .	11
1.2	Aerosols and climate . . . . .	11
1.3	Formation of secondary organic aerosol . . . . .	13
1.4	Partitioning coefficient . . . . .	15
1.5	Objectives of this work . . . . .	16
<b>2</b>	<b>Instrument description</b>	<b>17</b>
2.1	Proton Transfer Reaction-Time of Flight-Mass Spectrometer (PTR-ToF-MS) . . . . .	17
2.1.1	Working principle . . . . .	17
2.1.2	Ionization . . . . .	19
2.1.3	Mass resolution . . . . .	20
2.1.4	Data Handling . . . . .	22
2.2	Aerosol Collection Module (ACM) . . . . .	24
2.2.1	Working principle . . . . .	25
2.2.2	ACM collection section . . . . .	26
<b>3</b>	<b>Experiments</b>	<b>33</b>
3.1	AIDA chamber . . . . .	33
3.2	Instrumental set-ups and performances . . . . .	35
3.2.1	Set-up of the PTR-ToF-MS and characterization of the working conditions . . . . .	36
3.2.2	Set-up of the ACM and characterization of the working conditions . . . . .	38
3.3	Experimental Conditions and Procedure . . . . .	40
3.3.1	$\beta$ -pinene ozonolysis experiments . . . . .	41
3.3.2	$\alpha$ -pinene ozonolysis experiments . . . . .	44
3.3.3	Nopinone experiment . . . . .	45

## CONTENTS

---

<b>4 Procedures of data analysis</b>	<b>47</b>
4.1 Determination of the mixing ratios . . . . .	47
4.2 Calibration . . . . .	50
4.3 Fragmentation . . . . .	54
4.3.1 $\beta$ -Pinene Fragmentation . . . . .	57
4.3.2 $\alpha$ -Pinene Fragmentation . . . . .	59
4.3.3 Comparison of $\beta$ -pinene and $\alpha$ -pinene fragmentation to literature . . . . .	60
4.3.4 Nopinone fragmentation . . . . .	61
4.4 Determination of the partitioning coefficient from a PTR-ToF- MS in a filter set-up . . . . .	62
4.5 Determination of the partitioning coefficient from a PTR-ToF- MS coupled with the ACM . . . . .	65
<b>5 Results and discussion of <math>\beta</math>- and <math>\alpha</math>-pinene ozonolysis experi-     ments</b>	<b>67</b>
5.1 $\alpha$ -Pinene Ozonolysis . . . . .	67
5.1.1 $\alpha$ -Pinene Ozonolysis Products . . . . .	70
5.2 $\beta$ -Pinene Ozonolysis . . . . .	73
5.3 $\beta$ -Pinene Ozonolysis Products . . . . .	78
5.3.1 Nopinone Partitioning . . . . .	78
5.3.2 Other $\beta$ -Pinene Ozonolysis Products . . . . .	89
5.4 ACM - SMPS comparison . . . . .	91
5.5 Scientific findings of the $\alpha$ - and $\beta$ -pinene ozonolysis experiments	94
<b>6 Summary and Outlook</b>	<b>95</b>
<b>A Detailed descriptions of the experiment processes</b>	<b>99</b>
<b>B Pearson correlation matrix for <math>\beta</math>-pinene fragmentation</b>	<b>107</b>
<b>C Ion identities and Pearson correlation coefficient for ions cor-     related to <math>C_{10}H_{17}^+</math></b>	<b>109</b>
<b>D Mixing ratio error</b>	<b>113</b>
<b>E Fragments from ionization of <math>C_{10}H_{16}</math> with <math>O_2^+</math></b>	<b>115</b>
<b>F Ions in the aerosol particle phase of a <math>\beta</math>-pinene experiment     at 243 K</b>	<b>117</b>
<b>References</b>	<b>119</b>

## **CONTENTS**

---

<b>List of Figures</b>	<b>125</b>
<b>List of Tables</b>	<b>127</b>
<b>Abbreviations</b>	<b>129</b>
<b>Danksagung</b>	<b>131</b>



**Energie & Umwelt / Energy & Environment**  
**Band / Volume 169**  
**ISBN 978-3-89336-862-4**

