



Radiolytic Stability of BTBP-, BTPPhen- and DGA-based Ligands for the Selective Actinide Separation by Solvent Extraction

Holger Schmidt

Energie & Umwelt / Energy & Environment

Band / Volume 617

ISBN 978-3-95806-723-3

Forschungszentrum Jülich GmbH
Institut für Energie- und Klimaforschung (IEK)
Nukleare Entsorgung (IEK-6)

Radiolytic Stability of BTBP-, BTPhen- and DGA-based Ligands for the Selective Actinide Separation by Solvent Extraction

Holger Schmidt

Schriften des Forschungszentrums Jülich
Reihe Energie & Umwelt / Energy & Environment

Band / Volume 617

ISSN 1866-1793

ISBN 978-3-95806-723-3

Table of Contents

Abstract.....	I
Kurzzusammenfassung	III
List of Abbreviations	VII
1. Introduction.....	1
1.1. Back-end of the nuclear fuel cycle.....	2
1.2. Hydrometallurgical reprocessing: solvent extraction.....	4
2. Radiolysis in nuclear reprocessing.....	11
2.1. Radiolysis of the aqueous phase	11
2.2. Radiolysis of the organic phase	14
2.2.1. Radiolysis of alcohols	15
3. Ligands of interest for this work	17
3.1. Nitrogen donor ligands.....	17
3.1.1. Development and evolution of N-donor ligands	17
3.1.2. Radiolytic degradation of BTBP-type ligands	20
3.1.3. Pulse radiolysis of BT(B)Ps.....	21
3.2. Diglycolamides (DGA).....	22
3.2.1. Development and evolution of Diglycolamides.....	22
3.2.2. Radiolysis of Diglycolamides	23
4. Scope of the work and objectives	27
5. Materials and Methods.....	29
5.1. Experiments using N-donor extractants.....	29
5.2. Experiments using DGA extractants.....	31
6. Conclusions.....	33

6.1. Summary of the work on CyMe ₄ BTBP and CyMe ₄ BTPPhen	33
6.2. Summary of the work on DGA-type ligands	38
7. List of Figures	43
8. References.....	45
9. Acknowledgements.....	59
10. List of Papers	61

Energie & Umwelt / Energy & Environment
Band / Volume 617
ISBN 978-3-95806-723-3